

Arthroscopic Rotator Cuff Repair Small to Medium Tears With Type II SLAP repair

Precautions: No Active Biceps for 8 weeks
No CKC exercises for 8 weeks
No excessive IR for 4 weeks
No excessive horizontal adduction for 4 weeks

I. **Phase I - Immediate Post-Surgical Phase (Days 1-10)**

Goals: Maintain Integrity of the Repair
Gradually Increase Passive Range of Motion
Diminish Pain and Inflammation
Prevent Muscular Inhibition

Days One to Six:

- Abduction pillow brace
- Pendulum Exercises
- Active Assisted ROM Exercise (L-Bar)
 - ER/IR in Scapular Plane at 45 degrees of abduction (pain-free ROM)
- Passive ROM
 - Flexion to tolerance (painful ROM)
 - ER/IR in Scapular Plane at 45 degrees of abduction (pain-free ROM)
- Elbow/Hand Gripping & ROM Exercises
- Submaximal Painfree Isometrics (initiate days 4-5)
 - Flexion with elbow bent to 90 degrees
 - External Rotation
 - Internal Rotation
 - Elbow Flexors

- Cryotherapy for Pain and Inflammation
 - Ice 15-20 minutes every hour

- Sleeping
 - Sleep in pillow brace

Days Seven to Ten:

- Continue use of pillow brace
 - Pendulum Exercises
 - Progress Passive ROM to Tolerance
 - Flexion to at least 115 degrees
 - ER in Scapular Plane at 45 degrees abduction to 20-25 degrees
 - IR in Scapular Plane at 45 degrees abduction to 30-35 degrees
 - Active Assisted ROM Exercises (L-bar)
 - ER/IR in Scapular Plane at 45 degrees abduction
 - Flexion to Tolerance*
- *Therapist Provides Assistance by Supporting Arm (esp. with arm lowering)
- Continue Elbow/Hand ROM & Gripping Exercises

- Continue Isometrics (submaximal and subpainful)
 - Flexion with Bent Elbow
 - Extension with Bent Elbow
 - Abduction with Bent Elbow
 - ER/IR with Arm in Scapular Plane
 - Elbow Flexion
- Initiate rhythmic stabilization ER/IR at 45 degrees abduction
- Continue Use of Ice for Pain Control
 - Use Ice at least 6-7 times daily
- Sleeping
 - Continue Sleeping in Brace until Physician Instructs

Precautions:

1. No Lifting of Objects
2. No Excessive Shoulder Extension
3. No Excessive Stretching or Sudden Movements
4. No Supporting of Body Weight by Hands
5. Keep Incision Clean & Dry

II. Phase II - Protection Phase (Day 15 – Week 6)

Goals: Allow Healing of Soft Tissue
Do Not Overstress Healing Tissue
Gradually Restore Full Passive ROM (Week 4-5)
Re-Establish Dynamic Shoulder Stability
Decrease Pain & Inflammation

Days 15 – 21:

- Continue Use of Sling or Brace (physician or therapist will determine when to discontinue)
 - Passive Range of Motion to Tolerance
 - Flexion to 140-155 degrees
 - ER at 90 degrees abduction to at least 45 degrees
 - IR at 90 degrees abduction to at least 45 degrees
 - Active Assisted ROM to Tolerance
 - Flexion (continue use of arm support)
 - ER/IR in Scapular Plane at 45 degrees abduction
 - ER/IR at 90 degrees Abduction
 - Dynamic Stabilization Drills
 - Rhythmic Stabilization Drills
 - ER/IR in Scapular Plane
 - Flexion/Extension at 100 degrees Flexion and 125 degrees flexion
 - Continue All Isometric Contractions
 - Initiate scapular isometrics
 - Continue Use of Cryotherapy as needed
 - Continue All Precautions
 - No lifting
 - No excessive motion
-

Weeks 4 - 5:

- Patient should exhibit full passive range of motion by week 4
- Continue all exercises listed above
- Initiate ER/IR strengthening using exercise tubing at 0 degrees of abduction (use towel roll)
- Initiate Manual Resistance ER Supine in Scapular Plane (light resistance)
- Initiate Prone Rowing to Neutral arm Position
- Initiate prone shoulder extension
- Initiate ER strengthening exercises
- Initiate Isotonic Elbow Flexion
- Continue use of ice as needed
- May use heat prior to ROM exercises
- May use pool for light AROM exercises
- Rhythmic stabilization exercises (flexion 45, 90, 125 degrees) (ER/IR)

Weeks 5 – 6:

- May use heat prior to exercises
- Continue AAROM and Stretching exercises
 - Especially for movements that are not full
 - Shoulder flexion
 - ER at 90 degrees abduction
- Initiate Active ROM Exercises
 - Shoulder Flexion Scapular Plane
 - Shoulder Abduction
- Progress Isotonic Strengthening Exercise Program
 - ER Tubing
 - Sidelying IR
 - Prone Rowing
 - Prone Horizontal Abduction (bent elbow)

Precautions:

1. No Heavy Lifting of Objects
2. No excessive behind the back movements
3. No Supporting of Body Weight by Hands & Arms
4. No Sudden Jerking Motions

III. Phase III – Intermediate Phase (Weeks 7-14)

Goals: Full Active ROM (Week 8-10)
 Maintain Full Passive ROM
 Dynamic Shoulder Stability
 Gradual Restoration of Shoulder Strength
 Gradual Return to Functional Activities

Week 7:

- Continue Stretching & PROM (as needed to maintain full ROM)
- Continue Dynamic Stabilization Drills
- Passive Elbow Flexion only
- Progress Strengthening Program
 - ER/IR Tubing
 - ER Sidelying
 - Lateral Raises*

- Full Can in Scapular Plane*
- Prone Rowing
- Prone Horizontal Abduction
- Prone Extension
- Elbow Extension

*Patient must be able to elevate arm without shoulder or scapular hiking before initiating isotonic; if unable, continue glenohumeral joint exercises.

Week 8:

- Continue all exercise listed above
- If physician permits, may initiate Light functional activities

Week 10:

- Continue all exercise listed above
- Progress to Fundamental Shoulder Exercises
- Therapist may initiate isotonic resistance (1 lb wt.) during flexion and abduction*
 - *If non-painful normal motion is exhibited!

Weeks 11-15:

- Progress all exercises
 - Continue ROM and flexibility exercises
 - Progress strengthening program (increase 1 lb/10 days *non-painful)

IV. Phase IV – Advanced Strengthening Phase (Weeks 16 - 22)

Goals: Maintain Full Non-Painful ROM
Enhance Functional Use of UE
Improve Muscular Strengthen & Power
Gradual Return to Functional Activities

Week 16:

- Continue ROM & Stretching to maintain full ROM
- Self Capsular Stretches
- Progress Shoulder Strengthening Exercises
 - Fundamental Shoulder Exercises
- Initiate Interval Golf Program (if appropriate)

Weeks 20- 22:

- Continue all exercises listed above
- Progress Golf Program to playing golf (if appropriate)
- Initiate Interval Tennis Program (if appropriate)
- May Initiate Swimming

V. Phase V – Return to Activity Phase (Weeks 23 - 36)

Goals: Gradual Return to Strenuous Work Activities
Gradual Return to Recreational Sport Activities

Week 23:

- Continue Fundamental Shoulder Exercise Program (at least 4 times weekly)
 - Continue Stretching, if motion is tight
 - Continue Progression to Sport Participation
-